



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12Q	A2	(11) International Publication Number: WO 99/11813 (43) International Publication Date: 11 March 1999 (11.03.99)
<p>(21) International Application Number: PCT/US98/18397</p> <p>(22) International Filing Date: 3 September 1998 (03.09.98)</p> <p>(30) Priority Data: 60/057,810 4 September 1997 (04.09.97) US</p> <p>(71) Applicant: CHIRON DIAGNOSTICS CORPORATION [US/US]; 333 Coney Street, East Walpole, MA 02032 (US).</p> <p>(72) Inventors: HORN, Thomas; Chiron Diagnostics Corporation, 333 Coney Street, East Walpole, MA 02032 (US). SCHROEDER, Hartmut, R.; Chiron Diagnostics Corporation, 333 Coney Street, East Walpole, MA 02032 (US). WARNER, Brian, D.; Chiron Diagnostics Corporation, 333 Coney Street, East Walpole, MA 02032 (US). FISS, Ellen; Chiron Diagnostics Corporation, 333 Coney Street, East Walpole, MA 02032 (US). SELLS, Todd; Chiron Diagnostics Corporation, 333 Coney Street, East Walpole, MA 02032 (US). LAW, Say-Jong; Chiron Diagnostics Corporation, 333 Coney Street, East Walpole, MA 02032 (US).</p> <p>(74) Agent: MORGENSTERN, Arthur, S.; Chiron Diagnostics Corporation, 63 North Street, Medfield, MA 02052 (US).</p>		<p>(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> <p>Published <i>Without international search report and to be republished upon receipt of that report.</i></p>
<p>(54) Title: OLIGONUCLEOTIDE PROBES BEARING QUENCHABLE FLUORESCENT LABELS, AND METHODS OF USE THEREOF</p> <p>(57) Abstract</p> <p>Methods are provided for reducing background signals encountered in nucleic acid hybridization assays and other assays that involve hybridization of a labeled oligomer to its complement. The method is premised on the significant reduction of signal generation that occurs when a quenchable dye-labeled oligomer forms a hybrid complex. In addition, a method is provided for enhancing the detectable signal emitted from an amplification multimer hybridized to an oligomer probe to which a quenchable dye has been conjugated through a linker such that the emission from the dye is not quenched upon hybrid complex formation. Novel oligonucleotide probes are also provided that comprise an oligomer to which has been directly or indirectly through a linker a quenchable dye.</p> <div data-bbox="1036 1182 1377 1503"> </div>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

OLIGONUCLEOTIDE PROBES BEARING
QUENCHABLE FLUORESCENT LABELS,
AND METHODS OF USE THEREOF

5

10

TECHNICAL FIELD

This invention relates generally to nucleic acid chemistry and
15 hybridization assays. More particularly, the invention relates to a method for
detecting a target nucleic acid in a sample using an oligonucleotide probe
bearing a detectable label.

20

BACKGROUND

Nucleic acids have joined antigens and antibodies as key molecular
targets for human diagnostic assays. In particular, due to the recent
development of a variety of technical capabilities, microorganisms can now be
25 detected and quantified at unprecedented low levels in clinical specimens.

A variety of concepts have provided the basis for a number of different probe-target hybridization detection systems.

5 Glazer et al. (1992) *Nature* 359:959 describe fluorescent intercalation complexes of polycationic ligands with double-stranded DNA (dsDNA) for high-sensitivity DNA detection. The fluorescent homodimers intercalate into dsDNA to form a complex that is stable to electrophoresis, thus allowing detection and quantitation of DNA after separation with minimum background interference.

10 Tyagi et al. (1996) *Nature Biotech.* 14:303-308 describe a probe that is optically silent in solution but fluoresce upon hybridization with a complementary target. The probe is a single-stranded oligonucleotide that possesses a stem-and-loop structure. The loop is a sequence complementary to the target. One arm of the stem has a fluorescent moiety attached and the other arm has a nonfluorescent quenching moiety attached. In the nonhybridized state, the stem keeps the two moieties in sufficiently close proximity so that the fluorescent moiety is quenched by the nonfluorescent moiety. Upon hybridization, the consonant conformational change in the probe forces the arm sequences apart and allows the fluorescent moiety to be detected.

15 Mergny et al. (1994) *Nucleic Acids Res.* 22:920-928 describes the use of fluorescence energy transfer to probe for primary or secondary structural features of single-stranded DNA. Two probes that hybridize to adjacent sequence of the target DNA respectively carry donor and acceptor dyes. Hybridization of the two probes to the target leads to fluorescence excitation energy transfer between the donor and acceptor dyes.

20 Davis et al. (1996) *Nucleic Acids Res.* 24:702-706 describe the use of DNA constructs containing one or two fluorescein molecules in flow cytometry. The fluorescein molecules were attached to the 3'-end of a DNA probe through an 18-atom spacer arm that resulted in a 10-fold increase in fluorescence intensity compared to the DNA probe to which fluorescein was directly attached to the 3'-end of the probe.

25

30

In addition, many forms of target amplification have been introduced for enhancing detection sensitivity since the polymerase chain reaction was first developed. These methods can be used for producing greater quantities of the target nucleic acid and assays using these methods generally use conventional
5 detection schemes.

In contrast, direct analysis of target nucleic acids with great sensitivity has been accomplished using signal amplification. This method has been applied to the quantitation of many organisms and mRNAs. As few as 50 picomoles of the human immunodeficiency virus genome has been quantified in
10 human plasma samples. The key molecule in the amplification method is a branched DNA (bDNA) that permits the specific incorporation of many labeled probes.

Two types of bDNA molecules have been used. The first type is a large network comprised of DNA oligomers assembled in solution from
15 oligonucleotides containing three equally distributed N⁴-[N-(6-aminocaproyl-2-aminoethyl)]-5-methyl-2'-deoxycytidine residues coupled to one another through homobifunctional amine-reactive cross-linking agents. Such "amplification multimer" (AM) networks permitted the specific hybridization of many alkaline phosphatase-labeled DNA probes per hybridization probe-target
20 nucleic acid complex. These AMs gave significant signal amplification. As the multimer was increased in size, a limit to the signal amplification was achieved, possibly due to the location of complementary oligonucleotide sequences buried in the spherical cross-linked structure which may be inaccessible to the large alkaline phosphatase-labeled probes.

25 The second type of bDNA are "comb-type" molecules that comprise a linear primary oligonucleotide with a plurality of secondary side sequences having sites complementary to labeled DNA probes. As with the AMs, alkaline phosphatase-labeled DNA probes provide an amplified signal that does not match the theoretical signal amplification, possibly due to steric considerations.

The signal from fluorescently labeled DNA-probes can also be amplified using multiply labeled oligomers. However, due possibly to quenching effects, the incorporation of additional labels into a labeled probe does not produce a linear increase in fluorescent signal. For example, a two-fold increase in the number of labels per probe only yields a 1.2-fold increase in signal output. It would be possible to obtain an amplified fluorescent signal using a bDNA molecule as a scaffold for labeled DNA probes. However, again proximity of fluorophors may result in quenching-limited signal output. A highly sensitive fluorescent bDNA signal amplification-based assay would result if the quenching phenomenon were minimized.

SUMMARY OF THE INVENTION

The present invention provides methods and probes for detecting nucleic acid analytes in a sample. In general, the methods represent nucleic acid hybridization assays, such as fluorescent *in situ* hybridization assays, polymerase chain reaction assays, ligase chain reaction assays, competitive hybridization assays, strand displacement assays, and the like. In particular, the methods represent solution phase sandwich hybridization assays which involve binding the analyte to a solid support, labeling the analyte, and detecting the presence of label on the support. Preferred methods involve the use of amplification multimers which enable the binding of significantly more label in the analyte-probe complex, enhancing assay sensitivity and specificity. The probes comprise an oligonucleotide sequence to which has been conjugated, directly or indirectly through a linker, a dye that when in solution emits detectable radiation. Upon hybridization of the direct oligonucleotide-dye conjugate to a complementary oligonucleotide sequence the detectable radiation is substantially quenched. The detectable signal from an oligonucleotide-linker-

dye conjugate is not appreciably quenched upon hybridization to a complementary oligonucleotide.

In a first aspect of the invention, a method is provide for detecting an oligonucleotide of interest in a sample. The method comprises (a) providing an
5 oligonucleotide probe comprising (i) a nucleic acid sequence complementary to a nucleic acid sequence in the oligonucleotide of interest, and (ii) a label that, when the probe is in single-stranded, nonhybridized form, provides a detectable fluorescent signal, but which, when the probe hybridizes to a complementary nucleic acid strand, does not fluoresce, (b) combining the probe with the sample
10 suspected of containing the oligonucleotide of interest under hybridizing conditions, while monitoring emitted fluorescence from the probe, and (c) correlating any decrease in fluorescence which occurs throughout step (b) with the presence or quantity of the oligonucleotide of interest.

In a related aspect of the invention, an improved solution phase
15 sandwich hybridization assay for detecting an nucleic acid analyte in a sample, comprising: (a) binding the analyte indirectly to a solid support; (b) labeling the analyte; and (c) detecting the presence of label on the support, in which the improvement comprises incorporating a label probe system comprising (i) a label extender molecule having a first segment L-1 capable of hybridizing to a
20 nucleic acid sequence in the analyte and a second segment L-2, (ii) an amplification multimer containing a nucleic acid sequence M-1 capable of hybridizing to nucleic acid sequence L-2 and a plurality of identical oligonucleotide subunits containing nucleic acid sequences M-2 capable of hybridizing to a label probe, and (iii) an oligonucleotide probe comprising a
25 nucleic acid sequence L-3 capable of hybridizing to M-2 and a label coupled to the probe through a linker incapable of specifically hybridizing with a nucleic acid sequence in the analyte, the label extender or the amplification multimer.

The invention additionally encompasses an oligonucleotide probe comprising (i) a nucleic acid sequence complementary to a nucleic acid
30 sequence in an oligonucleotide of interest, and (ii) a label that, when the probe

is in single-stranded, nonhybridized form, provides a detectable fluorescent signal, but which, when the probe hybridizes to a complementary nucleic acid strand, does not fluoresce.

5 In addition, the invention encompasses a singly labeled oligonucleotide probe comprising (i) a single-stranded nucleic acid molecule comprising first and second complementary nucleotide sequences flanking a third nucleotide sequence that forms a loop structure when the first and second complementary nucleotide sequences hybridize with one another, wherein the third nucleotide sequence in the loop structure comprises a nucleotide sequence complementary
10 to a nucleotide sequence in a target oligonucleotide, and (ii) a label that, when the first and second nucleotide sequences are hybridized to one another, is substantially quenched, and when the third nucleotide sequence is hybridized to the oligonucleotide of interest and the first and second nucleotide sequences are in nonhybridized form, provides a detectable fluorescent signal.

15 These and other embodiments of the present invention will readily occur to those of ordinary skill in the art in view of the disclosure herein.

BRIEF DESCRIPTION OF THE FIGURES

20

FIG. 1 is a graphical representation of the results of the experiment described in Example 3. The triangles represent the fluorescence obtained for a quenchable dye-labeled probe free in solution. The diamonds represent the fluorescence obtained for a quenchable dye-labeled probe-amplification
25 multimer hybrid complex.

FIG. 2 is a graphical representation of the results of the experiment described in Example 3 in which a quenchable dye is conjugated to an oligomer probe through a T₃ linker. The triangles represent the fluorescence obtained for a quenchable dye-labeled probe free in solution. The diamonds represent the

fluorescence obtained for an oligomer-3'-T₅-quenchable dye-labeled probe-amplification multimer hybrid complex.

FIGS. 3A and 3B are illustrations depicting the use of doubly labeled fluorogenic probes in nick-translation PCR and in 5' nuclease assays, respectively, as described in Example 5. FIG. 3C is an illustration of the use of a probe labeled with a single quenchable dye species in a Taq polymerase-based assay system as described in Example 5.

FIGS. 4A and 4B illustrate the principle of operation of a doubly labeled molecular beacon and of a singly labeled hairpin probe, respectively, as described in Example 6.

FIG. 5 is an illustration depicting the detection and amplification of a single base change using the LCR fluorescence assay as described in Example 7.

FIG. 6 is a schematic representation of a strand displacement DNA probe assay as described in Example 9.

FIG. 7 is a schematic representation of a hybridization assay for detecting a point mutation as described in Example 10.

FIG. 8 is a schematic representation of a hybridization assay for detecting a point mutation as described in Example 11.

DETAILED DESCRIPTION OF THE INVENTION

Definitions and nomenclature:

Before the present invention is disclosed and described in detail, it is to be understood that this invention is not limited to specific assay formats, materials or reagents, as such may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only and is not intended to be limiting.

It must be noted that, as used in the specification and the appended claims, the singular forms "a," "an" and "the" include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to "a label probe" includes more than one label probe, reference to "a dye compound" includes more than one dye compound, reference to "a nucleic acid analyte" includes more than one such analyte, reference to "a nucleoside" includes more than nucleoside, and the like.

In this specification and in the claims which follow, reference will be made to a number of terms which shall be defined to have the following meanings:

The term "quench" or "quenching" is used to indicate a reduction in detectable emission radiation, e.g., fluorescent or luminescent radiation, from a source that would otherwise have emitted this radiation. Quenching is a reduction of at least 50%, preferably 80% and more preferably 90%, of the detectable radiation from the source.

The term "quenchable dye" as used herein is a single molecular species that emits detectable radiation when in solution or bound to a single-stranded oligomer, either directly or through a linking moiety. The detectable radiation of a quenchable dye bound directly to a single-stranded oligomer is reversibly quenched upon hybridization of the oligomer to a complementary oligonucleotide to form a hybrid duplex or triplex. No additional molecular species, e.g., a quenching dye, is required for the quenching to occur. However, if the quenchable dye is bound to the oligomer through a linker moiety, hybridization of the oligomer to its complement will not result in quenching of the detectable radiation emitted by the dye.

As used herein, the terms "polynucleotide" and "oligonucleotide" shall be generic to polydeoxyribonucleotides (containing 2-deoxy-D-ribose), to polyribonucleotides (containing D-ribose), to any other type of polynucleotide which is an N- or C-glycoside of a purine or pyrimidine base, and to other polymers containing nonnucleotidic backbones, for example, polyamide (e.g.,

peptide nucleic acids (PNAs)) and polymorpholino (commercially available from the Anti-Virals, Inc., Corvallis, Oregon, as Neugene™ polymers), and other synthetic sequence-specific nucleic acid polymers providing that the polymers contain nucleobases in a configuration which allows for base pairing and base stacking, such as is found in DNA and RNA. There is no intended distinction in length between the term "polynucleotide" and "oligonucleotide," and these terms will be used interchangeably. These terms refer only to the primary structure of the molecule. Thus, these terms include, for example, 3'-deoxy-2',5'-DNA, oligodeoxyribonucleotide N3'-P5' phosphoramidates, 2'-O-alkyl-substituted RNA, double- and single-stranded DNA, as well as double- and single-stranded RNA, DNA:RNA hybrids, and hybrids between PNAs and DNA or RNA, and also include known types of modifications, for example, labels which are known in the art, methylation, "caps," substitution of one or more of the naturally occurring nucleotides with an analog, internucleotide modifications such as, for example, those with uncharged linkages (e.g., methyl phosphonates, phosphotriesters, phosphoramidates, carbamates, etc.), with negatively charged linkages (e.g., phosphorothioates, phosphorodithioates, etc.), and with positively charged linkages (e.g., aminoalkylphosphoramidates, aminoalkylphosphotriesters), those containing pendant moieties, such as, for example, proteins (including nucleases, toxins, antibodies, signal peptides, poly-L-lysine, etc.), those with intercalators (e.g., acridine, psoralen, etc.), those containing chelators (e.g., metals, radioactive metals, boron, oxidative metals, etc.), those containing alkylators, those with modified linkages (e.g., alpha anomeric nucleic acids, etc.), as well as unmodified forms of the polynucleotide or oligonucleotide.

It will be appreciated that, as used herein, the terms "nucleoside" and "nucleotide" will include those moieties which contain not only the known purine and pyrimidine bases, but also other heterocyclic bases which have been modified. Such modifications include methylated purines or pyrimidines, acylated purines or pyrimidines, or other heterocycles. Modified nucleosides or

nucleotides will also include modifications on the sugar moiety, e.g., wherein one or more of the hydroxyl groups are replaced with halogen, aliphatic groups, or are functionalized as ethers, amines, or the like. The term "nucleotidic unit" is intended to encompass nucleosides and nucleotides.

5 Furthermore, modifications to nucleotidic units include rearranging, appending, substituting for or otherwise altering functional groups on the purine or pyrimidine base which form hydrogen bonds to a respective complementary pyrimidine or purine. The resultant modified nucleotidic unit may form a base pair with other such modified nucleotidic units but not with A, T, C, G or U.

10 Standard A-T and G-C base pairs form under conditions which allow the formation of hydrogen bonds between the N³-H and C⁴-oxy of thymidine and the N¹ and C⁶-NH₂, respectively, of adenosine and between the C²-oxy, N³ and C⁴-NH₂, of cytidine and the C²-NH₂, N¹-H and C⁶-oxy, respectively, of guanosine. Thus, for example, guanosine (2-amino-6-oxy-9-β-D-ribofuranosyl-

15 purine) may be modified to form *isoguanosine* (2-oxy-6-amino-9-β-D-ribofuranosyl-purine). Such modification results in a nucleoside base which will no longer effectively form a standard base pair with cytosine. However, modification of cytosine (1-β-D-ribofuranosyl-2-oxy-4-amino-pyrimidine) to form *isocytosine* (1-β-D-ribofuranosyl-2-amino-4-oxy-pyrimidine) results in a

20 modified nucleotide which will not effectively base pair with guanosine but will form a base pair with *isoguanosine*. *Isocytosine* is available from Sigma Chemical Co. (St. Louis, MO); *isocytidine* may be prepared by the method described by Switzer et al. (1993) *Biochemistry* 32:10489-10496 and references cited therein; 2'-deoxy-5-methyl-*isocytidine* may be prepared by the method of

25 Tor et al. (1993) *J. Am. Chem. Soc.* 115:4461-4467 and references cited therein; and *isoguanine* nucleotides may be prepared using the method described by Switzer et al. (1993), *supra*, and Mantsch et al. (1993) *Biochem.* 14:5593-5601, or by the method described U.S. Patent No. 5,780,610 to Collins et al. The nonnatural base pairs referred to as κ and π, may be

30 synthesized by the method described in Piccirilli et al. (1990) *Nature* 343:33-37

for the synthesis of 2,6-diaminopyrimidine and its complement (1-methyl-pyrazolo[4,3]pyrimidine-5,7-(4H,6H)-dione. Other such modified nucleotidic units which form unique base pairs have been described in Leach et al. (1992) *J. Am. Chem. Soc.* 114:3675-3683 and Switzer et al., *supra*, or will be
5 apparent to those of ordinary skill in the art.

The terms "polynucleotide analyte" and "nucleic acid analyte" are used interchangeably and refer to a single- or double-stranded nucleic acid molecule that contains a target nucleotide sequence. The analyte nucleic acids may be from a variety of sources, e.g., biological fluids or solids, food stuffs,
10 environmental materials, etc., and may be prepared for the hybridization analysis by a variety of means, e.g., proteinase K/SDS, chaotropic salts, or the like. The term "polynucleotide analyte" is used interchangeably herein with the terms "analyte," "analyte nucleic acid" and "target."

As used herein, the term "target region" or "target nucleotide sequence" refers to a probe-hybridizing region contained within the target molecule. The
15 term "target sequence" refers to a sequence with which a probe will form a stable hybrid under desired conditions.

As used herein, the term "probe" refers to a structure comprised of a polynucleotide, as defined above, that contains a nucleic acid sequence
20 complementary to a nucleic acid sequence present in the target analyte. The polynucleotide regions of probes may be composed of DNA, and/or RNA, and/or synthetic nucleotide analogs.

It will be appreciated that the hybridizing sequences need not have perfect complementarity to provide stable hybrids. In many situations, stable
25 hybrids will form where fewer than about 10% of the bases are mismatches, ignoring loops of four or more nucleotides. Accordingly, as used herein the term "complementary" refers to an oligonucleotide that forms a stable duplex with its "complement" under assay conditions, generally where there is about 90% or greater homology.

The terms "nucleic acid multimer" or "amplification multimer" are used herein to refer to a linear or branched polymer of the same repeating single-stranded oligonucleotide segment or different single-stranded polynucleotide segments, each of which contains a region where a labeled probe can hybridize, i.e., contains a nucleic acid sequence complementary to a nucleic acid sequence contained within a labeled probe; the oligonucleotide segments may be composed of RNA, DNA, modified nucleotides or combinations thereof. At least one of the segments has a sequence, length, and composition that permits it to hybridize specifically to a labeled probe; additionally, at least one of the segments has a sequence, length, and composition that permits it to hybridize specifically to a label extender or to a preamplifier system. Typically, such segments will contain approximately 15 to 50, preferably 15 to 30, nucleotides, and will have a GC content in the range of about 20% to about 80%. The total number of oligonucleotide segments in the multimer will usually be in the range of about 3 to 1000, more typically in the range of about 10 to 100, and most typically about 50. The oligonucleotide segments of the multimer may be covalently linked directly to each other through phosphodiester bonds or through interposed linking agents such as nucleic acid, amino acid, carbohydrate or polyol bridges, or through other cross-linking agents that are capable of cross-linking nucleic acid or modified nucleic acid strands. Alternatively, the multimer may be comprised of oligonucleotide segments which are in part covalently attached, but are also bonded in some other manner, e.g., through hybridization. Such a multimer is described, for example, in U.S. Patent No. 5,175,270 to Nilsen et al. The site(s) of linkage may be at the ends of the segment (in either normal, 3'-5' orientation or randomly oriented) and/or at one or more internal nucleotides in the strand. In linear multimers the individual segments are linked end-to-end to form a linear polymer. In one type of branched multimer three or more oligonucleotide segments emanate from a point of origin to form a branched structure. The point of origin may be another nucleotide segment or a multifunctional

molecule to which at least three segments can be covalently bound. In another type, there is an oligonucleotide segment backbone with one or more pendant oligonucleotide segments. These latter-type multimers are "fork-like," "comb-like" or combination "fork-" and "comb-like" in structure, wherein "comb-like" multimers, the preferred multimers herein, are polynucleotides having a linear backbone with a multiplicity of sidechains extending from the backbone. The pendant segments will normally depend from a modified nucleotide or other organic moiety having appropriate functional groups to which oligonucleotides may be conjugated or otherwise attached. The multimer may be totally linear, totally branched, or a combination of linear and branched portions. Typically, there will be at least two branch points in the multimer, more preferably at least three, more preferably in the range of about 5 to 30, although in some embodiments there may be more. The multimer may include one or more segments of double-stranded sequences. Further information concerning multimer synthesis and specific multimer structures may be found in commonly owned U.S. Patent No. 5,124,246 to Urdea et al.

Commonly assigned U.S. Patent Application Serial No. 07/813,588 and European Patent Publication No. 541,693 describe the comb-type branched multimers that are composed of a linear backbone and pendant sidechains; the backbone includes a segment that provides a specific hybridization site for analyte nucleic acid or nucleic acid bound to the analyte, whereas the pendant sidechains include iterations of a segment that provide specific hybridization sites for a labeled probe. A method of synthesizing a comb-type multimer is described below in the Experimental section.

As noted above, a "preamplifier system" may also be used, which serves as a bridging system between the label extender molecules and the amplification multimers. In this way, more amplifier and thus more label may be bound in any given target-probe complex. A preamplifier system contains a "prepreamplifier molecule" and/or a "preamplifier molecule." Prepreamplifier molecules are designed to serve as a bridge between label extender molecules

and preamplifier molecules, while preamplifier molecules are designed to serve as a bridge between either label extender molecules or prepreamplifier molecules and amplification multimers. Prepreamplifier molecules may be linear or branched, typically containing about 30 to about 3000 nucleotides, and
5 comprise at least one segment having a sequence, length, and composition that permits it to hybridize specifically to a label extender and at least one segment that permits it to hybridize to a preamplifier molecule. Preamplifier molecules, in turn, may be either linear or branched, and typically contain in the range of about 30 to about 3000 nucleotides. In one preferred embodiment herein, a
10 prepreamplifier or a preamplifier molecule hybridizes to at least two different label extender molecules, such that the overall accuracy of the assay is increased (i.e., because, again, a plurality of hybridization events are required for the probe-target complex to form).

As used herein, a "biological sample" refers to a sample of tissue or
15 fluid isolated from an individual, including but not limited to, for example, plasma, serum, spinal fluid, semen, lymph fluid, the external sections of the skin, respiratory, intestinal, and genitourinary tracts, tears, saliva, milk, blood cells, tumors, organs, and also samples of *in vitro* cell culture constituents (including but not limited to conditioned medium resulting from the growth of
20 cells in cell culture medium, putatively virally infected cells, recombinant cells, and cell components). Preferred uses of the present method are in detecting and/or quantitating nucleic acids as follows: (a) viral nucleic acids, such as from hepatitis B virus ("HBV"), hepatitis C virus ("HCV"), hepatitis D virus ("HDV"), human immunodeficiency virus ("HIV"), and the herpes family of
25 viruses, including herpes zoster (chicken pox), herpes simplex virus I & II, cytomegalovirus, Epstein-Barr virus, and the recently isolated Herpes VI virus; (b) bacterial nucleic acids, such as Chlamydia, Mycobacterium tuberculosis, etc.; and (c) numerous human sequences of interest.

"Label extender molecules (LEs)," also referred to herein as "label
30 extenders," contain regions of complementarity vis-à-vis the analyte

polynucleotide and to the amplification multimer. If a prepreamplifier or a preamplifier is used, the label extender molecules will hybridize to this intermediate species rather than directly to the amplification multimer. If neither a preamplifier system nor an amplifier is used, the label extender molecules will hybridize directly to a sequence in the labeled probe ("LP"). Thus, label extender molecules are single-stranded polynucleotide chains having a first nucleic acid sequence L-1 complementary to a sequence of the analyte polynucleotide, and a second universal region having a multimer recognition sequence L-2 complementary to a segment M-1 of label probe, amplification multimer, prepreamplifier or preamplifier.

"Labeled probes (LPs)" are designed to hybridize either to the label extender, or, if an amplification multimer is employed in the assay, to the repeating oligonucleotide segments of the multimer. LPs either contain a label or are structured so as to hybridize to a label. Thus, LPs contain a nucleic acid sequence L-3 complementary to a nucleic acid sequence M-2 present within the repeating oligonucleotide units of the multimer and are bound to, or structured so as to hybridize to, a label which provides, directly or indirectly, a detectable signal.

"Capture extender molecules (CEs)," also referred to herein as "capture extenders," hybridize to the analyte polynucleotide and to capture probes, which are in turn covalently bound to a solid support. Thus, capture extender molecules are single-stranded polynucleotide chains having a first polynucleotide sequence region containing a nucleic acid sequence C-1 which is complementary to a sequence of the analyte, and a second, noncomplementary region having a capture probe recognition sequence C-2. The sequences C-1 and L-1 are nonidentical, noncomplementary sequences that are each complementary to physically distinct sequences of the analyte.

"Capture probes (CPs)" hybridize to the capture extenders and bind to a solid support. Capture probes have a nucleic acid sequence C-3 complementary

to C-2 and are covalently bound to (or capable of being covalently bound to) a solid support.

Generally, solution phase hybridization capture assays are disclosed in U.S. Patent No. 5,635,352 to Urdea et al., the disclosure of which in
5 incorporated herein by reference, and proceed as follows. Single-stranded analyte nucleic acid is incubated under hybridization conditions with the capture extenders and label extenders. The resulting product is a nucleic acid complex of the analyte polynucleotide bound to the capture extenders and to the label
10 extenders. This complex may be subsequently added under hybridizing conditions to a solid phase having the capture probes bound to the surface thereof; however, in a preferred embodiment, the initial incubation is carried out in the presence of the support-bound capture probes. The resulting product comprises the complex bound to the solid phase via the capture extender
15 molecules and capture probes. The solid phase with bound complex is then separated from unbound materials. An amplification multimer, preferably a comb-type multimer as described above, is then optionally added to the solid phase-analyte-probe complex under hybridization conditions to permit the multimer to hybridize to the LEs; if a preamplifier system is used, the solid phase-analyte-probe complex is incubated with the preamplifier system probes
20 either along with the amplification multimer or, preferably, prior to incubation with the amplification multimer. The resulting solid phase complex is then separated from any unbound preamplifier and/or multimer by washing. The labeled probes are then added under conditions which permit hybridization to LEs, or, if an amplification multimer was used, to the repeating oligonucleotide
25 segments of the multimer. The resulting solid phase labeled nucleic acid complex is then washed to remove unbound labeled oligonucleotide. A label probe to which a quenchable dye has been conjugated will not emit a detectable signal while hybridized to an LE or an amplification multimer, as described more fully below. Thus, the label probe is then dissociated from the hybrid
30 complex with the LE or amplification multimer by any method well known in

the art, e.g., by warming, by the addition of a detergent or high salt, and the like. The signal from the dissociated label probe can then be read.


One primary focus of the present invention is on minimizing background noise by employing a probe that emits a detectable signal only under desired
5 conditions. Another primary focus of the invention is to provide a signal amplification system that produces a greater yield of detectable signal than possible with previous systems.

The inventors herein have observed that when an oligomer, directly labeled with a quenchable fluorescent dye, hybridizes to a complementary
10 nucleotide sequence, substantial quenching occurs relative to the fluorescence observed for the same amount of probe free in solution (*see* Example 1). As exemplified below, nucleic acid hybridization assays having reduced background noise by, for example, producing a detectable signal only upon dissociation of a directly labeled probe from its complement are a significant
15 improvement over conventional assay methods.

In addition, it has now been observed that quenching can be substantially reduced by coupling a quenchable dye to an oligomer through a linker (*see* Example 3). This provides a means to enhance signal amplification through the use of branched DNA amplification multimers. Conventional
20 fluorescent oligomer label probes having multiple labels per probe, tend to "self-quench," thereby limiting fluorescent signal emission. In addition, when a plurality of fluorescent label probes are hybridized to an amplification multimer containing a plurality of sequences complementary to the label probe, intramolecular quenching can occur. By contrast, hybridization of label probes
25 comprising a quenchable dye compound indirectly conjugated to an oligomer as disclosed and claimed herein to an amplification multimer does not appear to result in self-quenching and yields an increased signal output.

Preferred quenchable dyes are fluorescent dipyrrometheneboron difluoride dyes having the chemical structure

5



10 wherein R¹ through R⁷ are independently selected from the group consisting of:
hydrogen; halogen; alkyl; carboxyalkyl; acyl; aryl; arylalkyl; sulfonyl; formyl;
substituted ethenyl having the formula -CX=CY-Z wherein X, Y and Z are
independently halogen, C₁-C₁₀ alkyl, cyano, ester, amide, ethenyl, polyethenyl,
aryl or heteroaryl; and -L-G, wherein L is a linking moiety and G is a reactive
15 group enabling binding of the dye to an amino or carboxylic acid moiety;


or wherein (a) either R¹ and R², or R² and R³, and/or (b) either R⁴ and
R⁵, or R⁵ and R⁶, together form a benzene ring substituted with 0 to 4
substituents selected from the group consisting of hydrogen, halogen, cyano,
sulfonyl, sulfonate, carboxyl, carboxylate, alkyl, perfluoroalkyl, alkoxy,
20 alkylthio, nitro, amino, monoalkylamino, dialkylamino, alkylamido, aryl,
aryloxy, heteroaryl, heteroaryloxy, arylamino, heteroarylamino, or arylamido,
or wherein two *ortho* substituents are linked to form an additional such aromatic
ring,

with the proviso that at least one substituent within the group of R¹
25 through R⁷ is heteroaryl, and at least a second substituent within the group of R¹
through R⁷ is -L-G.

These compounds are described in U.S. Patent Nos. 4,774,339 to
Haugland et al., 5,187,288 to Kang et al., 5,248,782 to Haugland et al.,
5,274,113 to Kang et al., 5,451,663 to Kang et al., and 5,433,896 to Kang et
30 al., the disclosures of which are incorporated herein by reference. In

particular, U.S. Patent No. 5,541,663 describes fluorescent derivatives of
 dipyrrometheneboron difluoride dyes that are chemically reactive with
 biologically derived or chemically synthesized macromolecules to form dye-
 conjugates. Methods by which these fluorescent dyes may be synthesized and
 methods by which the dyes may be conjugated to biologically derived or
 synthetic chemical materials are also described in the Haugland et al. and Kang
 et al. patents cited herein. Many dipyrrometheneboron difluoride dyes useful in
 the invention are commercially available from Molecular Probes, Inc. (Eugene,
 OR).

A particularly preferred dipyrrometheneboron difluoride dye is



The dye may be conjugated directly to the oligomer or indirectly through a
 linker moiety. Oligomer-dye conjugates may be formed using conventional
 methods well known in the art.

The linker is a moiety that serves to connect the oligonucleotide probe to
 the label, covalently bound to both the probe and the label and having a
 molecular structure such that a desired degree of spacing is provided between
 the probe "backbone" and the label. Generally, the linker moiety is of a length
 sufficient to overcome the quenching effect of hybridization of the oligomer-
 label conjugate on the emission signal of the label. The linker may be a nucleic
 acid sequence, an amino acid sequence, a hydrocarbon chain, or the like.
 Preferably, the linker moiety is a nucleic acid sequence at least three or more
 nucleotides in length, more preferably between three and nine nucleotides in

length. In addition, it is preferred that the linker moiety not hybridize under standard hybridizing conditions to either (a) the nucleic acid sequence complementary to the label probe, or (b) the target nucleotide sequence. An additional preferred linker moiety is an ethylene glycol or triethylene glycol linker of three or more, preferably three to nine, ethylene glycol or triethylene glycol units. The linker moiety may be conjugated to the 5' or 3' end of the oligomer or to an internal nucleoside having suitably reactive side group.

In addition, the dye compound may be incorporated into an oligomer by covalent attachment to a derivatizable nucleotide. A modified nucleotide that has a derivatizable site to which a label may be attached, and oligomers containing such a modified nucleotide, are disclosed in U.S. Patent Nos. 4,910,300, 5,093,232, and 5,594,118 to Urdea et al., and 5,580,731 and 5,591,584 to Chang et al. For example, the N-4 modified pyrimidines of the Chang et al. '731 patent can be labeled at the N-4 position or at the 3' position.

As exemplified below, quenchable dye-oligomer conjugates are useful in a number of nucleic acid hybridization methods, including but not limited to solution-phase and solid-phase nucleic acid hybridization assays, *in situ* hybridization mapping, point mutation detection, and the like.

EXPERIMENTAL

The practice of the present invention will employ, unless otherwise indicated, conventional techniques of synthetic organic chemistry, biochemistry, molecular biology, and the like, which are within the skill of the art. Such techniques are explained fully in the literature. See, e.g., Sambrook, Fritsch & Maniatis, Molecular Cloning: A Laboratory Manual, Second Edition (1989); Oligonucleotide Synthesis (M.J. Gait, ed., 1984); Nucleic Acid

Hybridization (B.D. Hames & S.J. Higgins, eds., 1984); and the series, Methods in Enzymology (Academic Press, Inc.).

It is to be understood that while the invention has been described in conjunction with the preferred specific embodiments thereof, that the description above as well as the examples which follow are intended to illustrate and not limit the scope of the invention. Other aspects, advantages and modifications within the scope of the invention will be apparent to those skilled in the art to which the invention pertains.

In the following examples, efforts have been made to ensure accuracy with respect to numbers used (e.g., amounts, temperature, etc.) but some experimental error and deviation should be accounted for. Temperature is always given in degrees C and, unless otherwise indicated, pressure is at or near atmospheric.

General Methods

Polyacrylamide gel electrophoresis (PAGE) analysis: Native 10% polyacrylamide slab gels were prepared using a solution containing 10 ml of a 40% stock solution (Amresco, cat. no. 0496), 30 ml of 1x Tris-borate buffer ("TBE") (100 mM Tris-borate, 1 mM EDTA, 7 M urea, pH 8.3), 0.4 ml 10% ammonium persulfate and 20 μ l N,N,N',N'-tetramethylethylenediamine. A 0.5 mm slab was poured between 20 cm x 20 cm plates, and a comb with 10 1-cm teeth was inserted between the plates to form loading wells. After prerunning the gel at 25 mAmp for 20 min at 4°C and prewashing the loading wells, sample mixtures were loaded into the wells. The sample mixtures consisted of 5 μ l sample, 5 μ l 15% (w/v) Ficoll and 0.05% bromophenol blue (BPB) in water. The gel were run at 25 mAmp at 4°C. Bands were visualized by transferring the gel to a silica gel thin layer chromatography plate with a fluorescent indicator on the glass (Merck silica gel 60 F₂₅₄) which had been covered with plastic wrap. DNA bands were visualized by UV shadowing,


i.e., by exposing the gel on the plate to ultraviolet light at 260 nm. Fluorescent bands were visualized by exposing the gel on the plate to 360-nm light.

Oligonucleotide Synthesis: Oligodeoxynucleotides were synthesized by standard solid-phase chemistry using 2-cyanoethyl phosphoramidite monomers.

5 The phosphorylating reagent 2-((2-((4,4'-dimethoxytrityl)oxy)ethyl)sulfonyl)ethyl-2-cyanoethyl-N,N-diisopropylphosphoramidite (Phostel) was used to synthesize both 3'- and 5'-phosphorylated oligomers.

Conjugation of dye compounds to an oligonucleotide probe to form the label probe: The 3'-long chain amine ("LCA") portion (L) of the *bla*3
10 oligonucleotide (5'-AAG TAC GAC AAC CAC ATC L-3'), wherein L is N⁴-(6-aminocaproyl-2-aminoethyl)-5-methyl-2'-deoxycytidine, was reacted with the carboxylated dye molecule in a 1:100 ratio of BLA3:dye. Thus, the dye N-hydroxysuccinimide ester and BLA3 were added to 100 mM phosphate buffer,
15 pH 7.8, and incubated for 3 hr at room temperature.

The reaction mixture was applied to a NAP-5 column (Pharmacia) previously equilibrated with water and eluted with water. The desired product is isolated by PAGE. The product band is cut from the gel, crushed and extracted using 0.5 M NaCl, 100 mM Tris, pH 8.3, 10 mM MgCl₂, desalted
20 and ethanol precipitated. A BODIPY® FL LCA-nucleoside has the following structure:



If desired, the oligomer-dye conjugate may be further purified using, for example, ion exchange chromatography, hydrophobic chromatography, reverse phase chromatography, chromatofocusing or affinity chromatography. The ratio of label to DNA is determined using analytical gel electrophoresis.

5

EXAMPLE 1

Fluorescence Quenching Upon Hybridization of a Labeled Probe to a Linear Oligonucleotide

10

The functional properties of branched DNA amplification multimers were examined with two fluorescence techniques: a solution phase hybridization-dependent quenching test and a gel shift polyacrylamide analytical test.

15

A series of reaction mixtures each having a final volume of 5 μ l in 1x SSC was made containing 0-100 pmoles of each of the following components: (1) the amplification multimer prepared as described in Example 1; (2) 5'-BLA3-3'-dye or dye-5'-BLA3; and (3) 5'-BLA3c-PSCPc-3'. BLA3 is the oligomer

20

5'-AAG TAC GAC AAC CAC ATC TTT TT-3' and BLA3c-PSCPc is the oligomer 5'-GAT GTG GTT GTC GTA CTT TTT CTC TTG GAA AGA AAG TGA AGT G-3', wherein complementary oligomer segments are underlined. The reaction mixture was heated in a waterbath at 60°C for 15 min, then cooled to room temperature. The samples were diluted with 5 μ l

25

15% (w/v) Ficoll/0.05% BPB in water then loaded into the wells of a 10% native PAGE gel (20 x 20 cm poured without urea). The gel was electrophoresed until the tracer dye (BPB) had migrated about 2/3 down the gel. The bands were visualized by UV shadowing. The gel was transferred to a 20 x 20 cm silica gel thin layer chromatography plate (E. Merck silica gel 60 F₂₅₄

with fluorescent indicator on glass) which had been covered with plastic wrap. The gel/plate was exposed to UV light at 260 nm to visualize the DNA bands, and at 360 nm to visualize fluorescent bands. The 15x3 amplification multimer and the 15x3 amplification multimer/5'-BLA3-T₃-3'-BODIPY® FL hybrids did not migrate into the gel, whereas the 5'-BLA3-T₃-3'-BODIPY® FL migrated slightly ahead of the BPB dye.

A series of probes was synthesized with sequences complementary to repeated 18-nucleic acid sequences in a 60-mer. Each probe was labeled with BODIPY® FL (Molecular Probes, Eugene, OR), fluorescein or Texas Red. The individual probes were hybridized to a linear complementary sequence and analyzed by native PAGE to determine the fluorescence properties of the labeled oligomers. In all cases quantitative hybrid formation was achieved as determined by the complete disappearance of free labeled probes, but the observed fluorescence intensity varied. Extensive quenching was observed with the BODIPY® FL-labeled oligomer, in contrast to the fluorescein- and Texas Red-labeled oligomers which did not show an noticeable quenching under the same hybridization conditions. The BODIPY® FL dye was thus sensitive to the environment of the labeled oligomer, and quenching was a measure of the proportion of the probe that had hybridized with its complement to form a DNA duplex.

When the BODIPY® FL was linked to the probe through a T-T-T-T-T (T₅) linker at the 3' end of the probe, essentially no quenching was observed upon hybridization of the probe to its complement.

EXAMPLE 2

Synthesis of a 15-site branched comb DNA (bDNA)
and Assembly into an Amplification Multimer

5 A branched comb DNA having 15 branch sites to which was attached secondary sequences having three 18-mer hybridization sites complementary to a detectable label probe was prepared as follows.

 A. Synthesis of the primary sequence. Controlled pore glass (CPG) solid support (2000 Å pore size) derivatized with 14.8 μmol/gm of
10 dimethyltrityl-thymidine through an N-succinyl-aminopropyl linker (40.5 mg, 0.6 μmol) was packed into a column. The primary sequence was synthesized on an automated DNA synthesizer (Model 380B, Applied Biosystems Division, Perkin Elmer, Foster City, CA). The detritylation step used two 7-second pulses of trichloroacetate/CH₂Cl₂ (3% v/v), each followed by a 4-second pause,
15 then the column was flushed. This process was repeated two more times. The phosphoramidite reagent (18 μmol of a 90 mM solution in acetonitrile) was used for each coupling of A, G, C, or T; however, for the branching monomer (N-4-(6-hydroxyhexyl)-5-methyl-2'-deoxycytosine) phosphoramidite 20 μmol of a 100 mM solution in acetonitrile was used for each coupling.

20 B. Removal of levulinate protecting group of the branching monomer. The CPG from the primary synthesis was attached to a 10 ml plastic syringe and rinsed with 10 ml acetic acid/pyridine (1:1, v/v). Approximately 11 ml of 0.5 M hydrazine hydrate in acetic acid/pyridine was periodically pushed through over a period of 90 minutes at room temperature. The solid support
25 was then rinsed with 10 ml of the acetic acid/pyridine, detached from the syringe, and followed by extensive rinsing with acetonitrile before brief drying under vacuum.

 C. Synthesis of the secondary sequence. All reagents used to synthesize the secondary sequence were the same as in the primary sequence;

however, their quantities differed. The detritylation process used three 8-second pulses of trichloroacetate solution, each followed by a 4-second pause, followed by a 20-second rinse (a 15-second pulse with a 5-second pause) of toluene/CH₂Cl₂ (1:1, v/v). This process was repeated two more times.

5 Nucleoside phosphoramidite (96 μ mol) was used for each coupling of A, C, G, or T. The coupling process consisted of an 8-second addition of activator, an 8-second addition of activator and phosphoramidite, followed by a 30-second wait. This was performed a total of 8 times. The capping and oxidation process each used two 10-second pulses of reagents separated by a 5-second
10 pause, and then followed by a 60-second wait step. If the 5' ends of the secondary sequence required phosphorylation, then they were coupled with Phostel phosphoramidite (107 μ mol each at 100 mM) with no capping between couplings.

D. Deprotection and Purification of the 15-site bDNA. The detritylated
15 bDNA was cleaved with 2 ml 30% NH₄OH for 45 min at room temperature. The supernatant was collected in a 4-ml glass vial which was capped and heated in a 60°C oven for at least 16 hours, then dried under a vacuum. The crude bDNA was purified on a 7%T,5%C polyacrylamide gel (20 x 40 x 1.5 cm) containing 7 M urea. The gel was run until the bromophenol blue dye migrated
20 to within 2-3 cm of the bottom. The product band was excised and soaked in a 100 mM Tris-HCl buffer, pH 8.0 with 0.5 M NaCl and 5 mM EDTA for at least 24 hours with agitation. The salt was removed by loading the bDNA solution onto a short C-18 column and rinsing it with water. The purified bDNA was eluted from the column with methanol/H₂O (1:1, v/v). After being
25 dried and reconstituted in water, the bDNA was precipitated from ethanol/0.3 M aqueous potassium acetate (3:1, v/v).

E. Enzymatic Assembly of 15x3 bDNA Amplification Multimer. The 15-site bDNA comb oligomer (1 nmole), containing the 5'-p-TGA CTG-3' secondary sequence, and linear DNA sequences 5'-(GAT GTG GTT GTC GTA
30 CTT)₃-GCG TAG-3' (60-mer, 18.75 nmole) were combined with the linear

DNA linker 5'-CAG TCA CTA CGC-3' (12-mer, 18.75 nmole) in a reaction tube in a total volume of 140 μ l, and 25 μ l ligation buffer (10x buffer: 500 mM Tris, pH 7.5, 100 mM MgCl₂, 20 mM spermidine) was added. The tube was heated to 95°C and then slowly cooled to room temperature. At this point ATP (5 μ l of 0.1 M), polyethylene glycol 8000 (70 μ l of a 50% solution) and T4-ligase (6.7 units/ μ l, 50 units total) were added. This reaction was incubated at room temperature overnight. Sodium chloride was added (16.5 μ l of 4 M) and ice-cold ethanol (800 μ l) was added. The mixture was kept at -20°C for 30 minutes and then centrifuged at 12,000 x g for 30 min. The supernatant was decanted and the precipitate was first gently dried *in vacuo* and then resuspended in water. The product was purified as described above for the 15-site bDNA comb oligomer.

15

EXAMPLE 3

Solution Phase Hybridization to a 15x3
Amplification Multimer: Effect of Interposing
a Linker Between the Probe and the Fluorophor

20

The number of hybridizable BLA3c 18-mer sequences in the amplification multimer containing 15 branches, each of which has three BLA3c sequences (a 15x3 bAM) was determined using BODIPY® FL hybridization-dependent quenching. Increasing amounts of BLA3-3'-BODIPY® FL (0 to 60 pmoles) were incubated with the 15x3 amplification multimer. After hybridization and dilution, the fluorescence was measured. The results of this experiment are shown in FIG. 1.

25

A standard curve for labeled probe alone was constructed and shown to have a linear response with increasing probe concentration. In contrast, when the same amounts of BLA3-3'-BODIPY® FL were hybridized to the 15x3

amplification multimer the resulting curve had two distinct slopes. At lower concentrations of labeled probe the fluorescence output was quenched by about 80% and showed a flat slope. On addition of more labeled probe the slope changed to parallel that of the free nonhybridized labeled probe, suggesting saturation of hybridizable sites. From the interception of the two linear regions it was determined that the 15x3 amplification multimer contained an average of 36 hybridizable sites, or 12 60-mer sequences.

The hybridization experiment was repeated substituting BLA3-T₅-3'-BODIPY® FL for the BLA3-3'-BODIPY® FL. T₅ indicates a nucleotide spacer of 5 lined thymidines to which the dye is linked to the 3'-end of the BLA3 oligomer. In contrast to the results obtained using the BLA3-3'-BODIPY® FL probe, quenching was not observed using the BLA3-T₅-3'-BODIPY® FL probe. The two curves, depicted in FIG. 2, relative fluorescence from hybridized and nonhybridized labeled probe, were essentially superimposable.

This observation further suggested that the 15x3 amplification multimer provides a scaffold that could accommodate hybridization of multiple labeled probes without internal quenching. This would be consistent with previous reports that have noted that energy transfer between fluorophors occurs when labels are placed within 9 to 18 base pairs of each other. The length and rigidity of the linkage between the oligomer and the label also affected quenching between chromophors. Morrison (1995) in Kricka (ed.), *Nonisotopic Probing, Blotting, and Sequencing* (Academic Press, NY), pp. 430-471. The 15x3 amplification multimer and the T₅-modified labeled probe formed hybrids that provided rigid spacing of more than 18 base pairs between dye molecules.

EXAMPLE 4

Solution Phase Hybridization to a 15x3
Amplification Multimer with PAGE Detection:
Effect of Interposing a Spacer Between the
Oligomer Probe and the Fluorescent Dye

5 The hybridization properties of BODIPY® FL conjugates probes were examined as described in Example 3 using native PAGE analysis, which does not destabilize oligonucleotide hybrids during electrophoresis. The 15x3
10 amplification multimer was allowed to hybridize with increasing amounts of BLA3-T₅-3'-BODIPY® FL. The free labeled probe migrated into the gel; the 15x3 amplification multimer was too large to enter the gel. The gel analysis showed that when the BODIPY® FL-labeled probe was fully hybridized to the amplification multimer, all fluorescence was retained in the loading well and
15 did not migrate into the gel. As the amount of probe was increased beyond saturation of all hybridizable sites on the 15x3 amplification multimer, excess fluorescent probe migrated into the gel. The break-through when free probe started to migrate into the gel corresponded to approximately 35 hybridizable sites. The specificity of the hybridization was further demonstrated when the
20 same modified labeled probe was incubated with a 15x3 amplification multimer that contained a noncomplementary sequence. All of the labeled probe was seen to migrate into the gel and none of the fluorescence was retained in the loading well.

25

EXAMPLE 5

Single Label-Quenching PCR
Using Fluorogenic Probes

5 Lee et al. (1993) *Nucleic Acids Res.* 21:3761-3766 describe the use of a doubly labeled probe to monitor the results of nick-translation PCR to discriminate between alleles. In this method, a probe labeled at 5'-end with fluorescent-phosphoramidite and at 3'-end with amino modifier C6 dT Linker (Glen) and 3'-phosphate. An oligomer probe is converted to a double-labeled
10 oligomer with 5'-6-carboxyfluorescein, or 5'-TET and 3'-6-carboxytetramethyl rhodamine, a quencher dye. The labeled probe is designed to anneal to a region of one strand of a PCR product downstream from one of the PCR primers. The 3' end of the probe is blocked from extension by the 3' phosphate. The probe is included with the other PCR reagents, e.g., dNTPs, and is degraded by the
15 5'-3' exonuclease activity inherent in Taq polymerase as the primer is extended in every cycle. Because the probe is doubly labeled, i.e., has a fluorophor at the 5' end and a quencher at the 3' end, the fluorescence of the indicator dye is quenched. An illustration of this method is provided in FIG. 3A. Probe digestion during PCR restores indicator dye fluorescence if and only if
20 nucleolytic cleavage occurs between the two dyes. The degradation of two probes of different sequence can be monitored simultaneously if the two indicator dyes are spectrally distinguishable.

 Rudert et al. (1997) *BioTechniques* 22:1140-1145 describes the use of similarly doubly labeled probes in 5' nuclease assays. A primer oligomer
25 containing a hair-pin sequence and an oligomer probe having a 5'-fluorophor and a 3'-quencher are used. The primer forms a spontaneous hair-pin at the 3' end and hybridizes to the complementary sequence within the fluorescent probe. Due to the presence of the quencher, the fluorophor is not detectable when the probe is hybridized to the primer. In addition, upon hybridization, there is a

two-nucleotide gap between the hair-pin primer and the probe. Taq polymerase extends the 3' end of the primer and cleaves the 5' end of the probe, liberating the reporter dye from the quenching effect of the 3' quenching dye. This system is illustrated in FIG. 3B.

5 In either of these methods by which the activity of a nucleotide polymerase may be monitored, an oligonucleotide singly labeled with BODIPY® FL or other quenchable dye as defined herein at the 5'-terminal nucleotide will hybridize to a target sequence and the fluorescence will be quenched without the use of a quencher dye. Taq polymerase will digest the
10 probe from the 5'-end and release a 5'-BODIPY® FL-nucleotide, thus overcoming the quenching and producing a detectable signal. This system is illustrated in FIG. 3C.

15

EXAMPLE 6

Single Label Quenching Molecular Beacon

A Molecular Beacon system reports the presence of specific nucleic acids in homogeneous solution. These probes undergo a spontaneous
20 fluorogenic conformational change when they hybridize to their target. Only perfectly complementary targets elicit this response, as hybridization does not occur when the target contains a mismatched nucleotide or a deletion. The probes can be used to monitor the synthesis of specific nucleic acids in real time. When used in nucleic acid amplification assays, gene detection is
25 homogeneous and sensitive, and can be carried out in a sealed tube. *See*, Tyagi et al. (1996) *Nature* 14:303-308.

Probes are single-stranded nucleic acid molecules that possess a stem-and-loop structure (*see* FIG. 4A). The loop portion of the molecule is a probe sequence that is complementary to a predetermined sequence in the target

nucleic acid. The stem is formed by annealing of the two complementary arm sequences that are on either side of the probe sequence, and the arm sequences are unrelated to the target sequence.

5 A fluorophor is attached to the end of one arm and a nonfluorescent quencher moiety is attached to the end of the other arm. The stem structure is used to keep the fluorophor and the quencher in close proximity to each other and to ensure efficient quenching of the fluorophor by fluorescence resonance energy transfer (FRET) in the isolated probe. When the probe hybridizes to a target molecule, it forms a hybrid that is longer and, hence, more stable than
10 the hybrid stem structure. The probe undergoes a spontaneous conformational change that forces the arm sequences apart and causes the fluorophor and the quencher to separate. Thus, FRET is eliminated upon hybridization of the probe with the target and the fluorophor can now emit a detectable fluorescent signal.

15 This same design can be used without the need for a separate quenching dye (*see* FIG. 4B). The hairpin probe is labeled with a quenchable dye at one terminus. When in the hairpin structure, the quenchable dye fluorescence is essentially completely quenched because the terminal nucleotide is part of a duplex. When the beacon hairpin duplex is broken and the long duplex with the target is formed the quenchable dye-labeled terminus is no longer in a duplex
20 and quenching of the dye fluorescence is overcome.

Free, unhybridized probes do not emit a detectable radiation and, accordingly, it is not necessary to separate probes hybridized to the target from unhybridized probes. Accordingly, single label quenching molecular beacons
25 can be used for the detection of nucleic acids in homogeneous assays and in living cells, as well as for real time monitoring of assays in which nucleic acids are being synthesized, e.g., polymerase chain reactions.

EXAMPLE 7

Ligase Chain Reaction

In this method, illustrated in FIG. 5, a probe containing the two
5 sequences PSCPc (5'-TTT CTC TTG GAA AGA AAG TGA AGT G-3') and
NA1 (a first nucleic acid sequence complementary to a first portion of a target
sequence TA1) is specifically linked to another probe containing the sequences
NA2 (a second nucleic acid sequence complementary to a second portion of a
10 target sequence TA2) and LLA2 (the label extender sequence in the Chiron
Quantiplex assay) when both probes are hybridized to a target. Oligonucleotide
products are exponentially amplified by thermal cycling of the ligation reaction
in the presence of a second set of adjacent oligonucleotides that are
complementary to the first and the target. After the amplification the ligated
probe product, PSCPc-NA1-NA2-LLA2, is captured on a solid substrate, e.g.,
15 a plate or a bead, to which is attached a oligomer containing the PSCP sequence
via the PSCPc tail. The presence of the LLA2 tail sequence makes it possible
to use a bDNA amplification assay with a label probe conjugated through a
linker to a quenchable dye to detect the ligation product. (*see* FIG. 5).

20

25

EXAMPLE 8

Quenchable Fluorescent Oligomers

5 Incorporation of quenchable dye-labeled nucleotide triphosphates during enzymatic DNA/RNA synthesis produces a strongly fluorescent oligomer. However, the dye will be quenched when the oligomer is in a double- or triple-stranded hybrid complex. Separation of the strands by, e.g., exposure to increased temperature, will result in a strong fluorescent signal from the incorporated labeled nucleotides. Such oligomers are useful in nucleic acid
10 hybridization assays.

EXAMPLE 9

Strand Displacement Assay

15 Ellwood et al. (1986) *Clin Chem.* 32:1631-1636 describe an assay system for distinguishing between a native oligonucleotide strand and one containing a point mutation. In this system, a long probe oligomer is hybridized to a shorter labeled oligomer (20-50 nucleotides). The hybrid is
20 mixed with the sample believed to contain the mutant strand; any target DNA present in the sample will displace the labeled oligomer. To quantitate the results, the displaced oligomer must be separated from the hybrids by a suitable technique. This simple technique is potentially of great value for quantitation of nucleic acid species, but it is important to have a simple, reliable method of
25 separation.

 Using a quenchable dye-labeled probe the Ellwood et al. system will work without separation, i.e., it will work as a homogeneous assay, since in the double- stranded DNA form the quenchable dye fluorescence is quenched, whereas after displacement it will be fluorescent (*see* FIG. 6).

EXAMPLE 10

Point Mutation Detection

5 Saiki et al. (1985) *Biotechnology* 3:1008-1012, which is incorporated herein by reference, describes a method for detecting a gene containing a point mutation. A short ^{32}P -labeled oligonucleotide complementary to a target sequence in the wild type gene is prepared and hybridized to a sample DNA, thereby creating a new restriction enzyme cleavage site. If the sample contains
10 a gene having a point mutation within the target sequence, the mutation results in, for example, a mismatch base pair upon formation of the labeled oligonucleotide-mutant gene hybrid complex. The presence of the mismatch base pair interferes with cleavage of the oligonucleotide-mutant gene hybrid duplex by the restriction enzyme. Any nonhybridized labeled oligonucleotide is
15 blocked by the addition of a mutant-specific oligomer, i.e., an oligomer that hybridizes to the labeled oligonucleotide to form a complex that does not have the restriction enzyme cleavage site. Addition of the appropriate restriction enzyme will cut only the wild type gene, releasing a short labeled single-stranded fragment of the oligomer into solution. Separation by electrophoresis
20 and autoradiography will reveal whether the label is on a short fragment or, as in the case of the mutant allele, still bound to the gene.

 Using a quenchable dye-labeled probe the method can be modified to be conducted entirely in solution phase, without requiring component separation. After hybridization of the probe to the wild type gene, the restriction digest will
25 release part of the labeled probe in a short duplex with a portion of the target. This duplex can be designed to be quite short and of low stability resulting in spontaneous strand separation. The quenching of the quenchable dye fluorescence in the duplex is overcome and a detectable signal will result. If the mutant gene is present, the detectable signal will be substantially reduced or

eliminated. Excess quenchable dye-labeled probe remains hybridized to the mutant-specific oligomer, thus the dye signal is quenched and undetectable (*see* FIG. 7).

5

EXAMPLE 11

Point Mutation Detection

A second method for detecting a point mutation in a gene is illustrated in FIG. 8. In this method, a first probe is prepared having a first nucleic acid sequence that is complementary to a native target nucleic acid sequence in a gene. The target nucleic acid sequence is chosen to be in close proximity or adjacent to, and either upstream (as illustrated) or downstream from, a sequence suspected of containing a point mutation. The first probe has a second nucleic acid sequence complementary to a nucleic acid sequence in a second probe. The second nucleic acid sequence of the first probe is covalently linked to a quenchable dye molecule. A second probe is provided having a third nucleic acid sequence that is complementary to either the wild type nucleic acid sequence in the gene or, optionally, the nucleic acid sequence in the gene harboring the point mutation, as shown in FIG. 8. The wild type or mutant nucleic acid sequence in the gene is downstream (as illustrated) or upstream from the native target nucleic acid sequence. The second probe has a fourth nucleic acid sequence that is complementary to the second nucleic acid sequence of the first probe.

The assay can be conducted using, for example, two alternative protocols. In a first alternative, the gene of interest is incubated with the first probe under hybridizing conditions to form a gene-first probe hybrid complex. The gene-first probe hybrid complex is then incubated with the second probe in which the third sequence is either complementary to the wild type sequence or

complementary to the mutant nucleic acid sequence in the gene of interest (*see* FIG. 8). If the gene contains a wild type sequence and the second probe is complementary to the mutant sequence, the gene-first probe hybrid complex will not hybridize to the second probe and the emission radiation from the quenchable dye compound will not be affected. If the gene contains a mutant sequence and the second probe is complementary thereto, a gene-first probe-second probe complex will be formed and the emission radiation from the dye will be substantially quenched. Thus, a decrease in signal emission indicates the presence of a point mutation.

In a second alternative, the assay can be conducted using a second probe having a third sequence that is complementary to the wild type sequence. In this case, a decrease in signal emission indicates the presence of the wild type sequence, but not when the gene contains a mutant nucleic acid sequence.

EXAMPLE 12

High Resolution Fiber Fluorescent In Situ Hybridization Optical Mapping

High Resolution Fiber Fluorescent In Situ Hybridization (FISH) Optical Mapping is an extension of the methodology developed in the laboratory of David Schwartz (*see, e.g.,* Schwartz et al. (1997) *Curr. Opin. Biotechnol.* 8:70-74, and references cited therein; Samad et al. (1995) *Genome Res.* 5:1-4, and references cited therein; Meng et al. (1995) *Nat. Genet.* 9:432-438; and Cai et al. (1995) *Proc. Natl. Acad. Sci. USA* 92:5164-5168).

DNA is attached to a silanized coverslip in an elongated configuration. Once bound and stretched, the DNA is available for hybridization. Similar methodologies have been reported, *e.g.,* Weier et al. (1995) *Human Molec. Genet.* 4:1903-1910, in which the ability to label stretched DNA molecules with

sequence-specific labels by hybridization with biotin- and digoxigenin-labeled probes was demonstrated. However, multiple layers of fluorescence-labeled antibodies or fluorescence-labeled avidin molecules were required to generate a detectable fluorescent signal. The experiments described in this Example demonstrate the utility of branched DNA to label DNA in a sequence-specific manner, thereby obviating the need for multiple layers to detect specific regions of DNA.

A. λ DNA was labeled by hybridization using probes and the protocol reported by Weier et al for confirmation purposes. The lengths of the DNA restriction fragments were compared with reported results.

B. A model system was constructed to detect the HIV gag/pol region within a larger molecule. HIV gag/pol DNA was inserted into λ GT10 phage at the unique *EcoRI* site. The two methods of detection (Weier et al. vs. bDNA) were compared.

In the first method, probes were synthesized using random primer protocols. The gag/pol region was hybridized with probes labeled with digoxigenin followed by detection with rhodamine-labeled anti-digoxigenin antibody and Texas red-labeled anti-sheep antibody. The λ DNA was hybridized with fluorescein-labeled probes followed by detection with anti-fluorescein antibody and fluorescein-labeled anti-mouse antibody in multiple layers.

In the second method, bDNA technology was used to label the HIV portion of the λ GT10 construct followed by counterstaining the λ DNA with oxazole yellow homodimer (YOYO; Glazer et al. (1992) *Nature* 359:359). The results of the hybridizations were analyzed using IP Labs System software.

Fiber FISH mapping of λ DNA: The 6 kb *HindIII* fragment of λ was mapped with anti-digoxigenin Texas Red antibody and appeared as red spots along the FITC-labeled backbone. Fiber FISH mapping was able to map the 6 kb fragment within 2 kb of the actual map position.

Fiber FISH mapping of GT10/HIV DNA: The 2.8 kb HIV fragment was mapped with anti-digoxigenin Texas Red antibody and appeared as red spots along the FITC backbone. Fiber FISH mapping was able to map the 2.8 kb fragment within 1 kb of the actual map position.

5 Fiber FISH bDNA mapping of GT10/HIV DNA: The HIV fragment was mapped with bDNA Texas Red-label probe and appear as red spots along the YOYO labeled backbone. Using bDNA the HIV fragment was mapped to within 1 kb of the predicted map position comparable to the fiber FISH mapping.

10 These results demonstrate the usefulness of bDNA in mapping small regions of DNA on a large backbone. Not only was the time to completion greatly shortened using bDNA (1 day or less) but the fluorescence signal using bDNA was considerably higher.

15

EXAMPLE 13

Use of bDNA Signal Amplification in Flow Cytometry

20 The flow cytometry assay uses reagents identical to those used in microwell-format bDNA assays with the exception of the solid phase and the labeled probe. In the flow cytometry technique, microparticles (latex beads) are derivatized with capture oligonucleotide. Following signal amplification with bDNA and a fluorescent reporter probe, the flow cytometer measures the
25 fluorescence of the beads in the assay mixture. Unbound reporter molecules or particles of any size other than that of the beads (about 3 μm) are not detected. After signal amplification, the fluorescence of beads that have captured a single target molecule is clearly separated from that of beads that have not captured a target.

Thus, novel methods for generating a target-dependent signal in nucleic acid hybridization assays and other methods that involve the hybridization of a labeled oligomer to its complement have been disclosed. In addition, a oligomer-dye conjugate has been disclosed. Although preferred embodiments of the subject invention have been described in some detail, it is to be understood that obvious variations can be made without departing from the spirit and the scope of the invention as defined by the appended claims.

10

15


20

25

WE CLAIM:

1. A method for detecting an oligonucleotide of interest in a sample, comprising:
 - 5 (a) providing a first oligonucleotide probe comprising (i) a first nucleic acid sequence complementary to a first nucleic acid sequence in the oligonucleotide of interest, and (ii) a label that, when the probe is in single-stranded, nonhybridized form, provides a detectable emission radiation that, when the probe hybridizes to a complementary nucleic acid strand, is
10 substantially quenched;
 - (b) combining the first oligonucleotide probe with the sample suspected of containing the oligonucleotide of interest under hybridizing conditions to form a first probe-oligonucleotide of interest hybrid complex, while monitoring emitted radiation from the first oligonucleotide probe; and
15 (c) correlating any change in emitted radiation that occurs throughout step (b) with the presence or quantity of the oligonucleotide of interest.
2. The method of claim 1, wherein the label comprises a fluorescent
20 dye.
3. The method of claim 2, wherein the dye is a dipyrrometheneboron difluoride compound.
4. The method of claim 3, wherein the dye has the chemical
25 structure

5




wherein R^1 through R^7 are independently selected from the group consisting of:
 10 hydrogen; halogen; alkyl; carboxyalkyl; acyl; aryl; arylalkyl; sulfonyl; formyl;
 substituted ethenyl having the formula $-CX=CY-Z$ wherein X, Y and Z are
 independently halogen, C_1-C_{10} alkyl, cyano, ester, amide, ethenyl, polyethenyl,
 aryl or heteroaryl; and $-L-G$, wherein L is a linking moiety and G is a reactive
 group enabling binding of the dye to an amino or carboxylic acid moiety;

15 or wherein (a) either R^1 and R^2 , or R^2 and R^3 , and/or (b) either R^4 and
 R^5 , or R^5 and R^6 , together form a benzene ring substituted with 0 to 4
 substituents selected from the group consisting of hydrogen, halogen, cyano,
 sulfonyl, sulfonate, carboxyl, carboxylate, alkyl, perfluoroalkyl, alkoxy,
 alkylthio, nitro, amino, monoalkylamino, dialkylamino, alkylamido, aryl,
 20 aryloxy, heteroaryl, heteroaryloxy, arylamino, heteroarylamino, or arylamido,
 or wherein two *ortho* substituents are linked to form an additional such aromatic
 ring,

with the proviso that at least one substituent within the group of R^1
 through R^7 is heteroaryl, and at least a second substituent within the group of R^1
 25 through R^7 is $-L-G$.

5. The method of claim 4, wherein the dye is

30



6. The method of claim 1, wherein the dye compound is coupled directly to the label probe.

10

7. The method of claim 1, wherein the dye compound is coupled indirectly to the label probe.

8. The method of claim 7, wherein the dye compound is coupled to the label probe through a linker substantially incapable of specifically hybridizing with a nucleic acid sequence in the oligonucleotide of interest.

15

9. The method of claim 8, wherein the linker is a nucleic acid sequence.

20

10. In a solution phase sandwich hybridization assay for detecting a nucleic acid analyte in a sample, comprising (a) binding the analyte indirectly to a solid support, (b) labeling the analyte, and (c) detecting the presence of label on the support, the improvement which comprises:

25 incorporating a label probe system comprising (i) a label extender molecule having a first segment L-1 capable of hybridizing to a nucleic acid sequence in the analyte and a second segment L-2, (ii) an amplification multimer containing a nucleic acid sequence M-1 capable of hybridizing to nucleic acid sequence L-2 and a plurality of identical oligonucleotide subunits containing nucleic acid sequences M-2 capable of hybridizing to a label probe,

30

(iii) a label probe comprising a nucleic acid sequence L-3 capable of hybridizing to M-2 and a quenchable dye coupled to the probe through a linker incapable of specifically hybridizing with a nucleic acid sequence in the analyte, the label extender, the amplification multimer or a target nucleotide sequence.

5


11. The assay of claim 10, wherein the quenchable dye comprises a fluorescent dye.

10

12. The assay of claim 11, wherein the dye is a dipyrrometheneboron difluoride dye.

13. The assay of claim 12, wherein the dye has the chemical structure

15



20

wherein R¹ through R⁷ are independently selected from the group consisting of: hydrogen; halogen; alkyl; carboxyalkyl; acyl; aryl; arylalkyl; sulfonyl; formyl; substituted ethenyl having the formula -CX=CY-Z wherein X, Y and Z are independently halogen, C₁-C₁₀ alkyl, cyano, ester, amide, ethenyl, polyethenyl, aryl or heteroaryl; and -L-G, wherein L is a linking moiety and G is a reactive group enabling binding of the dye to an amino or carboxylic acid moiety;

25


or wherein (a) either R¹ and R², or R² and R³, and/or (b) either R⁴ and R⁵, or R⁵ and R⁶, together form a benzene ring substituted with 0 to 4 substituents selected from the group consisting of hydrogen, halogen, cyano,

30

sulfonyl, sulfonate, carboxyl, carboxylate, alkyl, perfluoroalkyl, alkoxy, alkylthio, nitro, amino, monoalkylamino, dialkylamino, alkylamido, aryl, aryloxy, heteroaryl, heteroaryloxy, arylamino, heteroarylamino, or arylamido, or wherein two *ortho* substituents are linked to form an additional such aromatic ring,

with the proviso that at least one substituent within the group of R¹ through R⁷ is heteroaryl, and at least a second substituent within the group of R¹ through R⁷ is -L-G.

14. The assay of claim 13, wherein the dye is




15. The assay of claim 10, wherein the linker is a nucleic acid sequence.

16. An oligonucleotide probe comprising (i) a nucleic acid sequence complementary to a nucleic acid sequence in an oligonucleotide of interest, and (ii) a label that, when the probe is in single-stranded, nonhybridized form, provides a detectable fluorescent signal that, when the probe hybridizes to a complementary nucleic acid strand, is substantially quenched.

17. The oligonucleotide probe of claim 16, wherein the label comprises a fluorescent dye.

18. The oligonucleotide probe of claim 17, wherein the dye compound is a dipyrrometheneboron difluoride dye.

19. The oligonucleotide probe of claim 18, wherein the dye has the chemical structure




wherein R¹ through R⁷ are independently selected from the group consisting of:
hydrogen; halogen; alkyl; carboxyalkyl; acyl; aryl; arylalkyl; sulfonyl; formyl;
substituted ethenyl having the formula -CX=CY-Z wherein X, Y and Z are
independently halogen, C₁-C₁₀ alkyl, cyano, ester, amide, ethenyl, polyethenyl,
aryl or heteroaryl; and -L-G, wherein L is a linking moiety and G is a reactive
group enabling binding of the dye to an amino or carboxylic acid moiety;

or wherein (a) either R¹ and R², or R² and R³, and/or (b) either R⁴ and
R⁵, or R⁵ and R⁶, together form a benzene ring substituted with 0 to 4
substituents selected from the group consisting of hydrogen, halogen, cyano,
sulfonyl, sulfonate, carboxyl, carboxylate, alkyl, perfluoroalkyl, alkoxy,
alkylthio, nitro, amino, monoalkylamino, dialkylamino, alkylamido, aryl,
aryloxy, heteroaryl, heteroaryloxy, arylamino, heteroarylamino, or arylamido,
or wherein two *ortho* substituents are linked to form an additional such aromatic
ring,

with the proviso that at least one substituent within the group of R¹
through R⁷ is heteroaryl, and at least a second substituent within the group of R¹
through R⁷ is -L-G.

20. The oligonucleotide probe of claim 19, wherein the dye is



10 21. The oligonucleotide probe of claim 16, wherein the dye compound is coupled directly to the label probe.

22. The oligonucleotide probe of claim 16, wherein the dye compound is coupled indirectly to the label probe.

15

23. The oligonucleotide probe of claim 22, wherein the dye compound is coupled to the label probe through a linker substantially incapable of specifically hybridizing with a nucleic acid sequence in the oligonucleotide of interest.

20

24. The oligonucleotide probe of claim 23, wherein the linker is a nucleic acid sequence.

25. The method of claim 1, wherein:

25

the oligonucleotide of interest is a wild type gene and the first nucleic acid sequence in the first oligonucleotide probe is complementary to a target sequence in a wild type gene, wherein:

(i) formation of a hybrid duplex between the first oligonucleotide probe and the target sequence in the wild type gene creates a site cleavable by a specific restriction enzyme, and

30

(ii) formation of a hybrid duplex between the first oligonucleotide probe and a point mutation-containing target sequence in a variant of the wild type gene does not create a site cleavable by the restriction enzyme;

(iii) step (b) comprises


5 incubating the labeled first oligonucleotide probe with a sample suspected of containing the variant of the wild type gene to form a variant-probe hybrid;

adding to the sample an oligomer capable of hybridizing to the labeled first oligonucleotide probe to form a hybrid complex that does not
10 contain the restriction enzyme cleavage site; and

following the addition the oligomer, adding to the sample the restriction enzyme.

26. The method of claim 25, wherein the dye is a
15 dipyrrometheneboron difluoride compound.

27. The method of claim 26, wherein the dye has the chemical
structure




wherein R¹ through R⁷ are independently selected from the group consisting of:
hydrogen; halogen; alkyl; carboxyalkyl; acyl; aryl; arylalkyl; sulfonyl; formyl;
substituted ethenyl having the formula -CX=CY-Z wherein X, Y and Z are
30 independently halogen, C₁-C₁₀ alkyl, cyano, ester, amide, ethenyl, polyethenyl,

aryl or heteroaryl; and -L-G, wherein L is a linking moiety and G is a reactive group enabling binding of the dye to an amino or carboxylic acid moiety;

or wherein (a) either R¹ and R², or R² and R³, and/or (b) either R⁴ and R⁵, or R⁵ and R⁶, together form a benzene ring substituted with 0 to 4 substituents selected from the group consisting of hydrogen, halogen, cyano, sulfonyl, sulfonate, carboxyl, carboxylate, alkyl, perfluoroalkyl, alkoxy, alkylthio, nitro, amino, monoalkylamino, dialkylamino, alkylamido, aryl, aryloxy, heteroaryl, heteroaryloxy, arylamino, heteroarylamino, or arylamido, or wherein two *ortho* substituents are linked to form an additional such aromatic ring,

with the proviso that at least one substituent within the group of R¹ through R⁷ is heteroaryl, and at least a second substituent within the group of R¹ through R⁷ is -L-G.

28. The assay of claim 27, wherein the dye is



29. The method of claim 1, wherein:

the first oligonucleotide probe provided in step (a) is singly labeled at the 5'-terminal nucleotide;

step (a') comprises providing an oligonucleotide primer, wherein the primer is complementary to a second sequence in the oligonucleotide of interest and further wherein the first nucleic acid sequence in the oligonucleotide of

interest is downstream from the second nucleic acid sequence in the oligonucleotide of interest;


step (b) further comprises incubating the first oligonucleotide probe and the primer with a sample containing the oligonucleotide of interest under conditions in which the first oligonucleotide probe and the primer hybridize to the oligonucleotide of interest;

step (b') comprises adding an enzyme to the sample under conditions wherein the enzyme has 3' polymerase activity and 5' nuclease activity,

wherein the change in emitted radiation occurs in step (b') is an increase in emitted radiation.

30. The method of claim 29, wherein the dye is a dipyrrometheneboron difluoride compound.

31. The method of claim 30, wherein the dye has the chemical structure




wherein R¹ through R⁷ are independently selected from the group consisting of: hydrogen; halogen; alkyl; carboxyalkyl; acyl; aryl; arylalkyl; sulfonyl; formyl; substituted ethenyl having the formula -CX=CY-Z wherein X, Y and Z are independently halogen, C₁-C₁₀ alkyl, cyano, ester, amide, ethenyl, polyethenyl, aryl or heteroaryl; and -L-G, wherein L is a linking moiety and G is a reactive group enabling binding of the dye to an amino or carboxylic acid moiety;

or wherein (a) either R¹ and R², or R² and R³, and/or (b) either R⁴ and R⁵, or R⁵ and R⁶, together form a benzene ring substituted with 0 to 4 substituents selected from the group consisting of hydrogen, halogen, cyano, sulfonyl, sulfonate, carboxyl, carboxylate, alkyl, perfluoroalkyl, alkoxy, alkylthio, nitro, amino, monoalkylamino, dialkylamino, alkylamido, aryl, aryloxy, heteroaryl, heteroaryloxy, arylamino, heteroarylamino, or arylamido, or wherein two *ortho* substituents are linked to form an additional such aromatic ring,

with the proviso that at least one substituent within the group of R¹ through R⁷ is heteroaryl, and at least a second substituent within the group of R¹ through R⁷ is -L-G.

32. The assay of claim 31, wherein the dye is




33. A singly labeled oligonucleotide probe comprising (i) a single-stranded nucleic acid molecule comprising first and second complementary nucleotide sequences flanking a third nucleotide sequence that forms a loop structure when the first and second complementary nucleotide sequences hybridize with one another, wherein the third nucleotide sequence in the loop structure comprises a nucleotide sequence complementary to a nucleotide sequence in a target oligonucleotide, and (ii) a label that, when the first and second nucleotide sequences are hybridized to one another, is substantially quenched, and when the third nucleotide sequence is hybridized to the

oligonucleotide of interest and the first and second nucleotide sequences are in nonhybridized form, provides a detectable fluorescent signal.

34. The oligonucleotide probe of claim 33, wherein the label
5 comprises a fluorescent dye.

35. The oligonucleotide probe of claim 34, wherein the dye
compound is a dipyrrometheneboron difluoride dye.

36. The oligonucleotide probe of claim 35, wherein the dye has the
10 chemical structure



20 wherein R¹ through R⁷ are independently selected from the group consisting of:
hydrogen; halogen; alkyl; carboxyalkyl; acyl; aryl; arylalkyl; sulfonyl; formyl;
substituted ethenyl having the formula -CX=CY-Z wherein X, Y and Z are
independently halogen, C₁-C₁₀ alkyl, cyano, ester, amide, ethenyl, polyethenyl,
aryl or heteroaryl; and -L-G, wherein L is a linking moiety and G is a reactive
25 group enabling binding of the dye to an amino or carboxylic acid moiety;


or wherein (a) either R¹ and R², or R² and R³, and/or (b) either R⁴ and
R⁵, or R⁵ and R⁶, together form a benzene ring substituted with 0 to 4
substituents selected from the group consisting of hydrogen, halogen, cyano,
sulfonyl, sulfonate, carboxyl, carboxylate, alkyl, perfluoroalkyl, alkoxy,
30 alkylthio, nitro, amino, monoalkylamino, dialkylamino, alkylamido, aryl,

aryloxy, heteroaryl, heteroaryloxy, arylamino, heteroarylamino, or arylamido, or wherein two *ortho* substituents are linked to form an additional such aromatic ring,

with the proviso that at least one substituent within the group of R¹ through R⁷ is heteroaryl, and at least a second substituent within the group of R¹ through R⁷ is -L-G.

37. The oligonucleotide probe of claim 36, wherein the dye is

10



15

38. The method of claim 1, further wherein:

step (a') comprises providing a second oligonucleotide probe comprising a second nucleic acid sequence complementary to the first nucleic acid sequence in the oligonucleotide of interest;


20

step (b') comprises combining the first probe-oligonucleotide of interest hybrid complex formed in step (b) with the second oligonucleotide probe under conditions that the first oligonucleotide probe will be displaced from the first probe-oligonucleotide of interest complex, while monitoring emitted radiation from the first oligonucleotide probe.

25

39. The method of claim 38, wherein the dye is a dipyrrometheneboron difluoride compound.

40. The method of claim 39, wherein the dye has the chemical structure




wherein R^1 through R^7 are independently selected from the group consisting of: hydrogen; halogen; alkyl; carboxyalkyl; acyl; aryl; arylalkyl; sulfonyl; formyl; substituted ethenyl having the formula $-CX=CY-Z$ wherein X, Y and Z are independently halogen, C_1 - C_{10} alkyl, cyano, ester, amide, ethenyl, polyethenyl, aryl or heteroaryl; and $-L-G$, wherein L is a linking moiety and G is a reactive group enabling binding of the dye to an amino or carboxylic acid moiety;

or wherein (a) either R^1 and R^2 , or R^2 and R^3 , and/or (b) either R^4 and R^5 , or R^5 and R^6 , together form a benzene ring substituted with 0 to 4 substituents selected from the group consisting of hydrogen, halogen, cyano, sulfonyl, sulfonate, carboxyl, carboxylate, alkyl, perfluoroalkyl, alkoxy, alkylthio, nitro, amino, monoalkylamino, dialkylamino, alkylamido, aryl, aryloxy, heteroaryl, heteroaryloxy, arylamino, heteroarylamino, or arylamido, or wherein two *ortho* substituents are linked to form an additional such aromatic ring,

with the proviso that at least one substituent within the group of R^1 through R^7 is heteroaryl, and at least a second substituent within the group of R^1 through R^7 is $-L-G$.

41. The assay of claim 40, wherein the dye is



42. The method of claim 1, wherein the first oligonucleotide probe is
 10 singly labeled and comprises a single-stranded nucleic acid molecule comprising
 first and second complementary nucleic acid sequences flanking a third nucleic
 acid sequence that forms a loop structure when the first and second
 complementary nucleic acid sequences hybridize with one another to form as
 stem structure, wherein the third nucleic acid in the loop structure comprises a
 15 nucleic acid sequence complementary to the first nucleic acid sequence in the
 oligonucleotide of interest, and wherein the label is such that,

when the first and second nucleic acid sequences are hybridized to form
 the stem structure, the emitted radiation is substantially quenched, and


when the third nucleic acid sequence is hybridized to the oligonucleotide
 20 of interest and the first and second nucleic acid sequences are in nonhybridized
 form, provides a detectable emission radiation.

43. The method of claim 42, wherein the label comprises a
 25 fluorescent dye.

44. The method of claim 43, wherein the dye compound is a
 dipyrrometheneboron difluoride dye.

45. The method of claim 44, wherein the dye has the chemical
 30 structure

5




wherein R^1 through R^7 are independently selected from the group consisting of:
 10 hydrogen; halogen; alkyl; carboxyalkyl; acyl; aryl; arylalkyl; sulfonyl; formyl;
 substituted ethenyl having the formula $-CX=CY-Z$ wherein X, Y and Z are
 independently halogen, C_1-C_{10} alkyl, cyano, ester, amide, ethenyl, polyethenyl,
 aryl or heteroaryl; and $-L-G$, wherein L is a linking moiety and G is a reactive
 group enabling binding of the dye to an amino or carboxylic acid moiety;
 15 or wherein (a) either R^1 and R^2 , or R^2 and R^3 , and/or (b) either R^4 and
 R^5 , or R^5 and R^6 , together form a benzene ring substituted with 0 to 4
 substituents selected from the group consisting of hydrogen, halogen, cyano,
 sulfonyl, sulfonate, carboxyl, carboxylate, alkyl, perfluoroalkyl, alkoxy,
 alkylthio, nitro, amino, monoalkylamino, dialkylamino, alkylamido, aryl,
 20 aryloxy, heteroaryl, heteroaryloxy, arylamino, heteroarylamino, or arylamido,
 or wherein two *ortho* substituents are linked to form an additional such aromatic
 ring,

with the proviso that at least one substituent within the group of R^1
 through R^7 is heteroaryl, and at least a second substituent within the group of R^1
 25 through R^7 is $-L-G$.

46. The method of claim 45, wherein the dye is

30



47. The method of claim 1, wherein the first oligonucleotide probe further comprises a second nucleic acid sequence complementary to a third nucleic acid sequence in a second probe,

10

wherein the first nucleic acid sequence in the oligonucleotide of interest is adjacent to a second nucleic acid sequence in the oligonucleotide of interest, wherein the second nucleic acid sequence in the oligonucleotide of interest is either a wild type nucleic acid sequence or a nucleic acid sequence containing a point mutation, and wherein the method further comprises

15

(a') providing a second oligonucleotide probe having third and fourth nucleic acid sequences, wherein the third nucleic acid sequence is complementary to either the wild type or the mutant second nucleic acid sequence in the oligonucleotide of interest and the fourth nucleic acid sequence is complementary to the second nucleic acid sequence in the first oligonucleotide probe;

20

wherein step (b) further comprises incubating the second oligonucleotide probe with the first probe-oligonucleotide of interest complex and

wherein step (c) comprises correlating any change in emitted radiation that occurs throughout step (b) with the presence of the wild type sequence or the mutant sequence.


25

48. The method of claim 47, wherein the label comprises a fluorescent dye.

30

49. The method of claim 48, wherein the dye compound is a dipyrrometheneboron difluoride dye.

50. The method of claim 49, wherein the dye has the chemical structure




wherein R¹ through R⁷ are independently selected from the group consisting of:
hydrogen; halogen; alkyl; carboxyalkyl; acyl; aryl; arylalkyl; sulfonyl; formyl;
substituted ethenyl having the formula -CX=CY-Z wherein X, Y and Z are
independently halogen, C₁-C₁₀ alkyl, cyano, ester, amide, ethenyl, polyethenyl,
aryl or heteroaryl; and -L-G, wherein L is a linking moiety and G is a reactive
group enabling binding of the dye to an amino or carboxylic acid moiety;

or wherein (a) either R¹ and R², or R² and R³, and/or (b) either R⁴ and
R⁵, or R⁵ and R⁶, together form a benzene ring substituted with 0 to 4
substituents selected from the group consisting of hydrogen, halogen, cyano,
sulfonyl, sulfonate, carboxyl, carboxylate, alkyl, perfluoroalkyl, alkoxy,
alkylthio, nitro, amino, monoalkylamino, dialkylamino, alkylamido, aryl,
aryloxy, heteroaryl, heteroaryloxy, arylamino, heteroarylamino, or arylamido,
or wherein two *ortho* substituents are linked to form an additional such aromatic
ring,

with the proviso that at least one substituent within the group of R¹
through R⁷ is heteroaryl, and at least a second substituent within the group of R¹
through R⁷ is -L-G.

51. The method of claim 50, wherein the dye is



10

15

20

25

1/7




FIG. 1




FIG. 2

2/7




FIG. 3A




FIG. 3C

FIG. 3B

3/7




FIG. 4A




FIG. 4B

4/7




FIG. 5




FIG. 6

6/7




FIG. 7




FIG. 8



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : C07H 21/00, C21Q 1/68</p>	<p>A3</p>	<p>(11) International Publication Number: WO 99/11813</p> <p>(43) International Publication Date: 11 March 1999 (11.03.99)</p>
<p>(21) International Application Number: PCT/US98/18397</p> <p>(22) International Filing Date: 3 September 1998 (03.09.98)</p> <p>(30) Priority Data: 60/057,810 4 September 1997 (04.09.97) US</p> <p>(71) Applicant: CHIRON DIAGNOSTICS CORPORATION [US/US]; 333 Coney Street, East Walpole, MA 02032 (US).</p> <p>(72) Inventors: HORN, Thomas; Chiron Diagnostics Corporation, 333 Coney Street, East Walpole, MA 02032 (US). SCHROEDER, Hartmut, R.; Chiron Diagnostics Corporation, 333 Coney Street, East Walpole, MA 02032 (US). WARNER, Brian, D.; Chiron Diagnostics Corporation, 333 Coney Street, East Walpole, MA 02032 (US). FISS, Ellen; Chiron Diagnostics Corporation, 333 Coney Street, East Walpole, MA 02032 (US). SELLS, Todd; Chiron Diagnostics Corporation, 333 Coney Street, East Walpole, MA 02032 (US). LAW, Say-Jong; Chiron Diagnostics Corporation, 333 Coney Street, East Walpole, MA 02032 (US).</p> <p>(74) Agent: MORGENSTERN, Arthur, S.; Chiron Diagnostics Corporation, 63 North Street, Medfield, MA 02052 (US).</p>		<p>(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> <p>Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</p> <p>(88) Date of publication of the international search report: 6 May 1999 (06.05.99)</p>
<p>(54) Title: OLIGONUCLEOTIDE PROBES BEARING QUENCHABLE FLUORESCENT LABELS, AND METHODS OF USE THEREOF</p> <div data-bbox="219 1155 1404 1690"> <p>The diagram illustrates the mechanism of a quenchable fluorescent probe. It shows a DNA template with a PCR primer and a probe labeled with a dye (Dye) and a quencher (Quencher). The probe is designed to hybridize with the target sequence. When the probe hybridizes, the dye and quencher are in close proximity, leading to quenching of the dye's fluorescence. The diagram also shows the probe hybridizing to a target sequence, forming a hairpin structure, and the Taq Polymerase extending the probe, which results in the separation of the dye and quencher, allowing fluorescence to be detected.</p> </div> <p>(57) Abstract</p> <p>Methods are provided for reducing background signals encountered in nucleic acid hybridization assays and other assays that involve hybridization of a labeled oligomer to its complement. The method is premised on the significant reduction of signal generation that occurs when a quenchable dye-labeled oligomer forms a hybrid complex. In addition, a method is provided for enhancing the detectable signal emitted from an amplification multimer hybridized to an oligomer probe to which a quenchable dye has been conjugated through a linker such that the emission from the dye is not quenched upon hybrid complex formation. Novel oligonucleotide probes are also provided that comprise an oligomer to which has been directly or indirectly through a linker a quenchable dye.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/18397

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07H21/00 C12Q1/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07H C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	LEE L G ET AL: "DNA SEQUENCING WITH DYE-LABELED TERMINATORS AND T7 DNA POLYMERASE: EFFECT OF DYES AND DNTPS ON INCORPORATION OF DYE-TERMINATORS AND PROBABILITY ANALYSIS OF TERMINATION FRAGMENTS" NUCLEIC ACIDS RESEARCH, vol. 20, no. 10, 1992, pages 2471-2483, XP002045850	16-19, 22,23
Y	see page 2477, Fig 4g. ddC-tetramethylbodipy ; page 2478, Fig 5d; Page 2478 Fig 6e.	1-51
X	WO 93 09185 A (MOLECULAR PROBES INC) 13 May 1993	16-24
Y	see abstract; claim 1; examples 26,27	1-51
	-/--	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

24 February 1999

Date of mailing of the international search report

12/03/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Scott, J

INTERNATIONAL SEARCH REPORT

In International Application No

PCT/US 98/18397

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	MCGRATH J C ET AL: "Fluorescent ligands for the study of receptors" TRENDS IN PHARMACOLOGICAL SCIENCES, vol. 17, no. 11, November 1996, page 393-399 XP004071066 see the whole document ---	1-51
Y	US 5 248 782 A (HAUGLAND RICHARD P ET AL) 28 September 1993 see column 1, line 18 - line 27; claim 1 ---	1-51
X	WO 97 00967 A (BAYLOR COLLEGE MEDICINE) 9 January 1997 see the whole document ---	16-24
Y	see the whole document ---	1-51
X	WO 96 15270 A (PERKIN ELMER CORP) 23 May 1996 see abstract ---	1,2
Y	see abstract ---	1-51
X	EP 0 713 921 A (HOFFMANN LA ROCHE) 29 May 1996 see claim 1 ---	1,2
Y	see claim 1 ---	1-51
P,X	EP 0 808 829 A (NISSHIN SPINNING) 26 November 1997 see page 2, line 10 - line 19 see page 6, line 20 - line 30 see page 19, line 30 - line 50 see claims 1,6,10; examples 10,12-14 -----	1-51

INTERNATIONAL SEARCH REPORT

Information on patent family members

Int. l. Application No

PCT/US 98/18397

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9309185 A	13-05-1993	US 5274113 A AT 161871 T CA 2122627 A DE 69223985 D DE 69223985 T EP 0612336 A US 5451663 A US 5723218 A	28-12-1993 15-01-1998 13-05-1993 12-02-1998 16-07-1998 31-08-1994 19-09-1995 03-03-1998
US 5248782 A	28-09-1993	NONE	
WO 9700967 A	09-01-1997	US 5614386 A US 5861287 A US 5728529 A AU 699939 B AU 6288696 A CA 2225531 A EP 0833936 A	25-03-1997 19-01-1999 17-03-1998 17-12-1998 22-01-1997 09-01-1997 08-04-1998
WO 9615270 A	23-05-1996	US 5538848 A AU 695561 B AU 4283696 A CA 2201756 A EP 0792374 A JP 10510982 T US 5723591 A	23-07-1996 13-08-1998 06-06-1996 23-05-1996 03-09-1997 27-10-1998 03-03-1998
EP 0713921 A	29-05-1996	US 5571673 A AU 700885 B AU 3901095 A CA 2163388 A CN 1133888 A JP 8220092 A NZ 280506 A	05-11-1996 14-01-1999 30-05-1996 24-05-1996 23-10-1996 30-08-1996 27-07-1997
EP 0808829 A	26-11-1997	JP 10287870 A US 5856479 A	27-10-1998 05-01-1999